# Recombinant SARS-CoV-2 S-trimer Protein (His Tag)

Catalog Number: PKSR030489 1 Publications



Note: Centrifuge before opening to ensure complete recovery of vial contents.

# **Description**

**Synonyms** 2019-nCov S protein;2019-nCoV Spike glycoprotein;2019-nCoV S glycoprotein

**Species** SARS-CoV-2 HEK293 Cells **Expression Host Sequence** Cys15-Gln1208 QHD43416.1 Accession Calculated Molecular Weight 136.6 kDa 170-220 kDa Observed molecular weight

C-His Tag

**Bioactivity** Loaded Human ACE-2-Fc on Protein A Biosensor, can bind 2019-nCoV S-trimer

Protein-His with an affinity constant of 0.125 nM as determined in BLI assay.

### **Properties**

Purity > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles. Storage


**Shipping** This product is provided as liquid. It is shipped at frozen temperature with blue

ice/gel packs. Upon receipt, store it immediately at < - 20°C.

**Formulation** Supplied as a 0.2 µm filtered solution of PBS, pH7.4.

Reconstitution Not Applicable

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### **Background**

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, Oacetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# **Recombinant SARS-CoV-2 S-trimer Protein (His Tag)**

Catalog Number: PKSR030489 1 Publications



protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>