

## Recombinant Human DR6/TNFRSF21 Protein (His Tag)

Catalog No. PKSH033441

### Description

<b>Synonyms</b>	Tumor Necrosis Factor Receptor Superfamily Member 21, Death Receptor 6, CD358, TNFRSF21, DR6
<b>Species</b>	Human
<b>Expression_host</b>	HEK293 Cells
<b>Sequence</b>	Gln42-Leu350
<b>Accession</b>	NP_055267.1
<b>Mol_Mass</b>	34.6 kDa
<b>AP_Mol_Mass</b>	58 kDa
<b>Tag</b>	C-6His
<b>Bio_activity</b>	Testing in progress

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Tumor Necrosis Factor Receptor Superfamily Member 21 (TNFRSF21) is a type I transmembrane receptor that includes four extracellular cysteine-rich motifs and a cytoplasmic death domain. DR6 is highly expressed in heart; brain; placenta; pancreas; lymph node; thymus and prostate. DR6 may activate NF-kappa-B and JNK to promote apoptosis and T-cell differentiation. In addition; DR6 binds with N-APP; which is released by the deprivation of Trophic-factor. It triggers caspase activation and degeneration of both neuronal cell bodies (via caspase-3) and axons (via caspase-6). DR6 is also expressed on the tumor cell lines and can be induced by TNF-α.

## SDS-PAGE

