

Recombinant Mouse EpCAM Protein (His Tag)

Catalog No. PKSM041381

Description

Synonyms	17 1A; 323/A3; Adenocarcinoma associated antigen; Adenocarcinoma-associated antigen; Antigen identified by monoclonal AUA1; AUA1; CD326; CD326 antigen; Cell surface glycoprotein Trop 1; Cell surface glycoprotein Trop 2; Cell surface glycoprotein Trop-1; CO 17A; CO17 1A; CO17A; DIAR5; EGP 2; EGP; EGP2; EGP314; EGP40; Ep CAM; Ep-CAM; EPCAM; EPCAM; EpCAM1; Epithelial cell adhesion molecule; Epithelial Cell Adhesion Molecule Intracellular Domain (EpCAM-ICD); Epithelial cell surface antigen; Epithelial cellular adhesion molecule; Epithelial glycoprotein 1; Epithelial glycoprotein 314; Epithelial glycoprotein; ESA; GA733 1; GA733 2; GA733-2; gastrointestinal tumor-associated antigen 2; 35-KD glycoprotein; gp4; hEGP 2; hEGP314; HNPCC8; Human epithelial glycoprotein 2; KS 1/4 antigen; KS1/4; KSA; Ly74; Lymphocyte antigen 74; M1S 1; M1S2; M4S1; Major gastrointestinal tumor associated protein GA733 2; Major gastrointestinal tumor-associated protein GA733-2; mEGP314; Membrane component chromosome 4 surface marker (35kD glycoprotein); Membrane component; chromosome 4; surface marker 1; Membrane component; chromosome 4; surface marker; MIC18; MK 1; Protein 289A; TACD1; TACSTD1; TROP1; Tumor associated calcium signal transducer 1; Tumor associated calcium signal transducer 2 precursor; Tumor-associated calcium signal transducer 1
Species	Mouse
Expression_host	E.coli
Sequence	Glu24-Thr266
Accession	Q99JW5-1
Mol_Mass	27.65 kDa
AP_Mol_Mass	30 kDa
Tag	N-His

Properties

Purity	>90 % as determined by reducing SDS-PAGE.
Endotoxin	Not tested.
Storage	Samples are stable for up to twelve months from date of receipt at -70°C. Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Formulation	Lyophilized from sterile PBS, pH 7.4

Background

Epithelial cell adhesion molecule (EpCAM,CD326) is a type I transmembrane glycoprotein that functions as a homophilic, epithelial-specific intercellular cell-adhesion molecule. In addition to cell adhesion, EpCAM is also involved in cellular signaling, cell migration, proliferation, and differentiation. EpCAM is highly expressed on most carcinomas and therefore of potential use as a diagnostic and prognostic marker for a variety of carcinomas, and has become a therapeutic target.

SDS-PAGE

