

## Recombinant Rat Interleukin-2/IL-2 Protein (His Tag)

Catalog No. PKSR030460

### Description

<b>Synonyms</b>	Aldesleukin, IL 2, IL-2, IL2, IL2, Interleukin 2, Interleukin-2, Involved in regulation of T cell clonal expansion, Lymphokine, POIL2, T Cell Growth Factor, T-cell growth factor, TCGF
<b>Species</b>	Rat
<b>Expression_host</b>	HEK293 Cells
<b>Sequence</b>	Ala 21-Gln 155
<b>Accession</b>	P17108
<b>Mol_Mass</b>	16.3 kDa
<b>AP_Mol_Mass</b>	16 kDa
<b>Tag</b>	C-His
<b>Bio_activity</b>	Testing in progress

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Interleukin-2(IL-2) is a O-glycosylated four  $\alpha$ -helix bundle cytokine that has potent stimulatory activity for antigenactivated T cells. It is expressed by CD4+ and CD8+ T cells,  $\gamma\delta$  T cells, B cells, dendritic cells, and eosinophils. Mature rat IL-2 shares 66% and 73% amino acid sequence identity with human and mouse IL-2, respectively. The receptor for IL-2 consists of three subunits that are present on the cell surface in varying preformed complexes. IL-2 is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.

## SDS-PAGE

