

# Recombinant Human Eotaxin-3/CCL26 Protein (aa 24-94)



Catalog Number:PKSH033733

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	C-C Motif Chemokine 26;CC Chemokine IMAC;Eotaxin-3;Macrophage Inflammatory Protein 4-Alpha;MIP-4-Alpha;Small-Inducible Cytokine A26;Thymic Stroma Chemokine-1;TSC-1;CCL26;SCYA26
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Thr24-Leu94
<b>Accession</b>	Q9Y258
<b>Calculated Molecular Weight</b>	8.5 kDa
<b>Observed molecular weight</b>	13 kDa
<b>Tag</b>	None

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 1mM EDTA, 20% Glycerol, pH 9.0.
<b>Reconstitution</b>	Not Applicable

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Chemokine Ligand 26 protein (CCL26) is a novel small cytokine belonging to the CC chemokine family which is involved in immunoregulatory and inflammatory processes. CCL26 is constitutively expressed in thymus; but only transiently expressed in phytohemagglutinin-stimulated peripheral blood mononuclear cells. It specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. CCL26, along with Eotaxin-1 and Eotaxin-2; selectively activates the CC chemokine receptor 3 (CCR3). The Eotaxin-3-CCR3 interaction may play an important role in allergic diseases such as atopic dermatitis and bronchial asthma. The full-length cDNA for CCL26 encodes a protein of 94 amino acids with a putative signal peptide of either 23 or 26 amino acid residues. Both the 71 and 68 amino acid residue variants of recombinant CCL26 demonstrate equal potency in inducing chemotaxis of a human CCR3-transfected cell line. Unlike most other CC chemokines; CCL26 maps to human chromosome 7q11.2;

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human Eotaxin-3/CCL26 Protein (aa 24-94)

Catalog Number:PKSH033733



within 40 kilobases of the Eotaxin-2 loci. CCL26 and Eotaxin-2 are unique in that they are the only chemokines identified to date that map to chromosome 7.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017