

Recombinant Rhesus macaque IL-33/Interleukin-33 Protein (His Tag)

Catalog No. PKSQ050072

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-33;IL-33;Interleukin-1 Family Member 11;IL-1F11;Nuclear Factor From High Endothelial Venules;NF-HEV;IL33;C9orf26;IL1F11;NFHEV
Species	Rhesus macaque
Expression Host	E.coli
Sequence	Ser112-Ile270
Accession	F7DLG4
Calculated Molecular Weight	20.3 kDa
Observed molecular weight	21 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-33 (IL-33) belongs to the IL-1 superfamily. IL33 is highly expressed in endothelial venules found in tonsils, Peyer patches and mesenteric lymph nodes, but almost undetectable in placenta. IL33 induces the production of T

For Research Use Only

helper-2 (Th2)-associated cytokines. IL-33 is a cytokine that mediates its biological effects by binding to and signals through IL1RL1/ST2 and IL-1 Receptor Accessory Protein (IL1RAP), activating intracellular molecules in the NF- κ B and MAP kinase signaling pathways that drive production of type 2 cytokines from polarized Th2 cells.