Recombinant Rhesus macaque CD27/TNFRSF7 Protein (Fc

Tag)

Catalog Number: PKSQ050077



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		4.0	
Desc	7277	\mathbf{T}	m
I I A WELL I		171.0	

Synonyms CD27 Antigen; CD27L Receptor; T-Cell Activation Antigen CD27; T14; Tumor

Necrosis Factor Receptor Superfamily Member 7;CD27;CD27;TNFRSF7

SpeciesRhesus macaqueExpression HostHEK293 CellsSequenceThr21-Ile192AccessionF7BYS2Calculated Molecular Weight46.5 kDaObserved molecular weight60-90 kDaTagC-Fc

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH

8.5.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the print

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

CD27 antigen, also known as CD27L receptor, T-cell activation antigen CD27, T14, S152, Tp55, TNFRSF7 and Tumor necrosis factor receptor for superfamily member 7,belongs to the TNF-receptor superfamily. CD27 is a single-pass type I membrane protein and exists as a homodimer form, containing three TNFR-Cys repeats. CD27 transduces signals that lead to the activation of NF-KappaB and MAPK8/JNK. CD27 is involved in regulating B-cell activation and immunoglobulin synthesis, binding to the ligand CD70. TRAF2 and TRAF5 have been shown to mediate the signaling process of CD27. CD27-binding protein (SIVA), which is a proapoptotic protein, can bind to CD27 and is thought to play

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Rhesus macaque CD27/TNFRSF7 Protein (Fc Tag)



Catalog Number:PKSQ050077

a key role in the apoptosis. CD27 is required for generation and long-term maintenance of T cell immunity.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>