
Recombinant Rhesus macaque CD160 Protein (Fc Tag)

Catalog No. PKSQ050081

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	CD160 antigen;CD160
Species	Rhesus macaque
Expression Host	HEK293 Cells
Sequence	Met1-Leu158
Accession	G7MG20
Calculated Molecular Weight	45 kDa
Observed molecular weight	55-65 kDa
Tag	C-Fc
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD160 antigen is a cell membrane protein which contains one Ig-likeV-type (immunoglobulin-like) domain. CD160 is a GPI-anchored lymphocyte surface receptor in which expression is mostly restricted to the highly cytotoxic CD56(dim) CD16(+)peripheral blood NK subset. CD160 is a receptor showing broad specificity for both classical and non-classical MHC class I molecules. CD160 is expressed in spleen, peripheral blood, and small intestine. Expression of CD160 is

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

restricted to functional NK and T cytotoxic lymphocytes. CD160 acts as a co-activator receptor for CD3-induced proliferation of CD4+ CD160+ T cells isolated from inflammatory skin lesions. Activated NK lymphocytes release a soluble form of CD160 that functionally impairs the MHC-I-specific cytotoxic CD8(+) T lymphocyte responsiveness.

For Research Use Only