Catalog Number: PKSR030474



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Synonyms	S protein RBD;2019-nCoV S protein RBD		
Species	SARS-CoV-2		
Expression Host	HEK293 Cells		
Sequence	Asn331-Val524		
Accession	QHD43416.1		
Calculated Molecular Weight	47.9 kDa		
Observed molecular weight	50-60 kDa		
Tag	C-mFc		
Bioactivity	Immobilized Human ACE-2-His(Cat#PKSH032068) at 10µg/ml (100 µl/well) can bind Recombinant 2019-nCoV Spike Protein (RBD, mFc Tag) (Cat#PKSR030474). The ED50 of Recombinant 2019-nCoV Spike Protein (RBD, mFc Tag) (Cat#PKSR030474) is 62 ng/ml.		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.		
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.		
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20° C.		
Formulation	Supplied as a 0.2 μ m filtered solution of PBS, pH7.4.		
Reconstitution	Not Applicable		
Data			

Da 20 90 60	MK	R	
40		-	
30			
20			
14		Ŧ	

k| 1

> 95 % as determined by reducing SDS-PAGE.

Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com Catalog Number: PKSR030474



cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com