

# Recombinant Human CD32a/FCGR2A Protein (167 His, His&AVI Tag), Biotinylated

Catalog Number:PKSH030290

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Low affinity immunoglobulin gamma Fc region receptor II-a;IgG Fc receptor II-a;CDw32;Fc-gamma RII-a;Fc-gamma-RIIa;FcRII-a;CD32;FCGR2A;FCG2;FCGR2A1;IGFR2;CD32A;CDw32;Fc gamma RIIA;FCG2;FcGR;FCGR2
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Met 210
<b>Accession</b>	P12318-1
<b>Calculated Molecular Weight</b>	23.6 kDa
<b>Observed molecular weight</b>	33 kDa
<b>Tag</b>	C-His-Avi
<b>Bioactivity</b>	1. Immobilized human IgG2 at 10 µg/ml (100 µl/well) can bind biotinylated human CD32 with a linear ranger of 0.16-0.8 µg/ml. 2. Using the Octet RED System, the affinity constant (Kd) of CD32a-AVI-His + BirA bound to human IgG was 12nM. 3. Labeling ratio of biotin to protein: 0.1-0.6.

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Receptors for the Fc region of IgG (FcγR) are members of the Ig superfamily that function in the activation or inhibition

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human CD32a/FCGR2A Protein (167 His, His&AVI Tag), Biotinylated



Catalog Number:PKSH030290

---

of immune responses. Human Fc $\gamma$ Rs are divided into three classes designated Fc $\gamma$ RI (CD64), Fc $\gamma$ RII (CD32), and Fc $\gamma$ RIII (CD16), which generate multiple isoforms, are recognized. The activating- type receptor either has or associates non-covalently with an accessory subunit that has an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Fc $\gamma$ RI binds IgG with high affinity and functions during early immune responses, whereas Fc $\gamma$ RII and RIII are low affinity receptors that recognize IgG as aggregates surrounding multivalent antigens during late immune responses. Three genes for human Fc $\gamma$ RII (A, B, and C) and one for mouse (Fc $\gamma$ RIIB), encoding type I transmembrane proteins with ITAM motifs (Fc $\gamma$ RII A and C) or ITIM motifs (Fc $\gamma$ RIIB) in their cytoplasmic domains, have been identified. Human CD32, also known as Low affinity immunoglobulin  $\gamma$  Fc region receptor II-a, Fc $\gamma$ RII A or FCGR2A Protein, is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. Associated with an ITAM-bearing adapter subunit, FcR $\gamma$ , CD32a delivers an activating signal upon ligand binding, and results in the initiation of inflammatory responses including cytolysis, phagocytosis, degranulation, and cytokine production.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017