

## Recombinant Human cytomegalovirus (HCMV) Glycoprotein B / gB Protein (His Tag)

Catalog No. PKSV030208

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Glycoprotein B:gB
<b>Species</b>	CMV
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Arg 777-Val 907
<b>Accession</b>	AAA45920.1
<b>Calculated Molecular Weight</b>	93.0 kDa
<b>Observed molecular weight</b>	130-140 kDa
<b>Tag</b>	C-His

### Properties

<b>Purity</b>	> 70 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 70 % as determined by reducing SDS-PAGE.

### Background

Cytomegalovirus (CMV) (human herpesvirus 5) glycoprotein B, also referred as CMV gB or gB, which belongs to the herpesviridae glycoprotein B family. It is a 97-amino acid glycoprotein encoded by the ORF of UL55. Cytomegalovirus

### For Research Use Only

Glycoprotein B protein is the most abundant component of the envelope, a target of neutralizing antibodies with at least two defined neutralizing epitopes and an essential replication component. Cytomegalovirus Glycoprotein B protein plays important roles in HCMV entry, cell-cell spread of internal virions, and fusion of infected cells. In addition, Cytomegalovirus Glycoprotein B protein is one envelope protein capable of heparin binding. It forms a physical association with host cell annexin II independent of the presence of calcium.