

Recombinant Mouse sFRP2 Protein (His Tag)

Catalog Number:PKSM040904



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AI851596;Sdf5
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Cys 295
Accession	NP_033170.1
Calculated Molecular Weight	32.5 kDa
Observed molecular weight	36 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The Secreted frizzled-related protein (SFRP) family consists of five secreted glycoproteins in humans (SFRP1~5) that act as extracellular signaling ligands. Each SFRP is approximately 300 amino acids in length and contains a cysteine-rich domain (CRD) that shares 30-50% sequence homology with the CRD of Frizzled (Fz) receptors, a putative signal sequence, and a conserved hydrophilic carboxy-terminal domain. SFRPs are able to bind Wnt proteins and Fz receptors in the extracellular compartment. The interaction between SFRPs and Wnt proteins prevents the latter from binding the Fz receptors. The Wnt pathway plays a key role in embryonic development, cell differentiation and cell proliferation. sFRP2 is a member of the SFRP family acting as soluble modulators of Wnt signaling and contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins called FZ domain and a NTR domain. sFRP2 inhibits

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse sFRP2 Protein (His Tag)

Catalog Number:PKSM040904



hypoxia induced endothelial cell apoptosis and increases endothelial cell migration. It prevents mesoderm specification and maintains the cells in the undifferentiated state. SFRP2 is also a novel stimulator of angiogenesis that stimulates angiogenesis via a calcineurin/NFAT pathway, thus is regarded as a favorable target for the inhibition of angiogenesis in solid tumors. Mouse sFRP2 is highly expressed in the eye and is also detected in heart and lung at low level.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com