

Recombinant Human NFATC4 protein (His Tag)

Catalog No. PKSH500001

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	cytoplasmic 4, NF ATc4, NF-AT3, NF-ATc4, NFAC4_HUMAN, NFAT3, NFATc4, Nuclear factor of activated T cells cytoplasmic 4, Nuclear factor of activated T cells cytoplasmic calcineurin dependent 4, Nuclear factor of activated T-cells, T cell transcription factor NFAT3, T-cell transcription factor NFAT3
Species	Human
Expression Host	E.coli
Sequence	Ser401-Phe683
Accession	Q14934-1
Calculated Molecular Weight	31.5 kDa
Observed molecular weight	31-33 kDa
Tag	N-His
Bioactivity	Immunogen(E-AB-40328)

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4., 5% trehalose, 5% mannitol, 0.01% tween-80. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the print
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Ca²⁺-regulated transcription factor that is involved in several processes, including the development and function of the immune, cardiovascular, musculoskeletal, and nervous systems. Involved in T-cell activation, stimulating the transcription of cytokine genes, including that of IL2 and IL4. Along with NFATC3, involved in embryonic heart development. Involved in mitochondrial energy metabolism required for cardiac morphogenesis and function. Transactivates many genes involved in the cardiovascular system, including AGTR2, NPPB/BNP (in synergy with GATA4), NPPA/ANP/ANF and MYH7/beta-MHC. Involved in the regulation of adult hippocampal neurogenesis. Involved in BDNF-driven pro-survival signaling in hippocampal adult-born neurons. Involved in the formation of long-term spatial memory and long-term potentiation (By similarity). In cochlear nucleus neurons, may play a role in deafferentation-induced apoptosis during the developmental critical period, when auditory neurons depend on afferent input for survival. Binds to and activates the BACE1/Beta-secretase 1 promoter, hence may regulate the proteolytic processing of the amyloid precursor protein (APP) . Plays a role in adipocyte differentiation. May be involved in myoblast differentiation into myotubes. Binds the consensus DNA sequence 5'-GGAAAAT-3' (Probable). In the presence of CREBBP, activates TNF transcription. Binds to PPARG gene promoter and regulates its activity. Binds to PPARG and REG3G gene promoters.

For Research Use Only