A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD5L/hAIM protein (His tag)

Catalog No. PKSH033902

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD5 antigen-like, Apoptosis inhibitor expressed by macrophages, hAIM, CT-2, IgM-

associated peptide, SP-alpha, CD5L, API6, UNQ203/PRO229

Species Human

Expression Host HEK293 Cells
Sequence Met1-Gly347
Accession O43866
Calculated Molecular Weight 37.1 kDa
Observed molecular weight 42 kDa
Tag C-His

Bioactivity Testing in progress

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD5 Antigen-Like (CD5L) is a soluble protein that belongs to group B of the scavenger receptor cysteine-rich (SRCR) superfamily and contains three SRCR domains. CD5L is a secreted glycoprotein and expressed by macrophages presentin

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

lymphoid tissues. It binds to myelomonocytic and lymphoid cells and may play an important role in the regulation of the innate and adaptive immune systems. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gram-negative bacteria and the SRCR domain one of CD5L retains both the LPS and LTA binding activities. Furthermore, CD5L seems to play a role as an inhibitor of apoptosis.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com