

## Recombinant Human CD82 (N-Fc)

Catalog No. PKSH033916

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	CD82 antigenC33;CD82 molecule;CD82;IA4;IA4C33 antigen;Inducible membrane protein R2;KAI1;KAI1GR15;ST6;ST6tspan-27;Suppressor of tumorigenicity 6 protein;Tetraspanin-27;TSPAN27;Tspan-27;TSPAN274F9
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Gly103-Gln225
<b>Accession</b>	P27701
<b>Calculated Molecular Weight</b>	40.3 kDa
<b>Observed molecular weight</b>	40-60 kDa
<b>Tag</b>	N-Fc

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

CD82 antigen, also known as Kai-1, is a widely expressed palmitoylated molecule of the tetraspanin superfamily.

### For Research Use Only

KAI1/CD82 is localized on cell membrane and form interactions with other tetraspanins, integrins and chemokines which are respectively responsible for cell migration, adhesion and signaling. CD82/Kai-1 is a component of the promiscuous TIMP-1 interacting protein complex on the cell surface of human adenocarcinoma cells and gives insight into tumorigenic metastatic potential. CD82/Kai-1 suppresses EMT in prostate cancer cells adhered to fibronectin leading to reduced cell migration and invasiveness. CD82/Kai-1 function is important for muscle stem cell function in muscular disorders. Overexpression of CD82/Kai-1 suppresses growth, migration and invasion of oral cancer cells and may be considered as a potential therapeutic target in oral cancer.