

Recombinant Human Butyrophilin Subfamily 1 Member A1/BTN1A1 (C-6His-Avi) Biotinylated

Catalog No. PKSH033984

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Butyrophilin Subfamily 1 Member A1;BT;BTN1A1;BTN
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala27-Arg242
Accession	Q13410
Calculated Molecular Weight	26.8 kDa
Observed molecular weight	35-40 kDa
Tag	C-His-Avi

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Butyrophilin Subfamily 1 Member A1 (BTN1A1) is the major protein associated with fat droplets in the milk. It belongs to the immunoglobulin superfamily. BTN1A1 acts as a specific membrane-associated receptor for the association of

For Research Use Only

cytoplasmic droplets with the apical plasma membrane. It is localized to the major histocompatibility complex (MHC) class I region of 6p. It may have arisen relatively recently in evolution by the shuffling of exons between 2 ancestral gene families. It is shown that BTN1A1 inhibits the proliferation of CD4 and CD8 T-cells activated by anti-CD3 antibodies, T-cell metabolism and IL2 and IFNG secretion.