

Recombinant Human PVRIG (C-mFc)

Catalog No. PKSH034031

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	C7orf15;CD112R;PVRIG;transmembrane protein PVRIG;C7orf15MGC138295;MGC104322;MGC138297;MGC2463
Species	Human
Expression Host	HEK293 Cells
Sequence	Thr41-Asp171
Accession	Q6DKI7
Calculated Molecular Weight	40.6 kDa
Observed molecular weight	45-50 kDa
Tag	C-Fc
Bioactivity	Loaded Biotinylated Human Nectin-2-His on His Biosensor, can bind Human PVRIG with an affinity constant of 168.5n M as determined in BLI assay.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Human PVRIG (poliovirus receptor related immunoglobulin domain-containing protein), also known as CD112 receptor (CD112R), is an approximately 34 kDa single transmembrane protein in the poliovirus receptor-like protein (PVR) family. The extracellular domain sequence of human and mouse PVRIG have approximately 65% similarity. PVRIG functions as a cell surface receptor for Nectin-2/CD112, a cell surface protein that is widely expressed on antigen-presenting cells and tumor cells. Disrupting the PVRIG/Nectin-2 interaction enhances human T cell response, suggesting PVRIG is a novel checkpoint for human T cells. PVRIG may act as a coinhibitory receptor that suppresses T-cell receptor-mediated signals.

For Research Use Only