

Recombinant Human Ig lambda constant 2 (C-6His)

Catalog No. PKSH034054

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Ig lambda constant 2;Immunoglobulin Lambda Constant 2;Ig Lambda C Domain;IGLC;IGLC2;immunoglobulin lambda constant 2 (Kern-Oz-marker);MGC20392;MGC45681
Species	Human
Expression Host	HEK293 Cells
Sequence	Gly1-Ser106
Accession	P0DOY2
Calculated Molecular Weight	12.1 kDa
Observed molecular weight	16 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Immunoglobulin lambda constant region is constant region of immunoglobulin light chains. Immunoglobulins, also known

For Research Use Only

as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. IGLC2 is 105 amino acids (aa) in length. It contains one Ig-like domain (aa 7-100) and generates a disulfide bond with an IgH chain via Cys104.