

Recombinant Mouse TGFBR3 (C-6His)

Catalog No. PKSM041420

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	InBGCAN;TBRIII;TGF-beta receptor type 3;TGF-beta receptor type III;TGF-beta RIII;TGFbetaRIII;TGFBR3;TGF-bRIII;TGFR-3;Transforming growth factor beta receptor III
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Glu23-Thr785
Accession	A0A0R4J097
Calculated Molecular Weight	85.9 kDa
Observed molecular weight	110-125 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Transforming growth factor beta receptor III (TGF-beta RIII; also betaglycan) is a ubiquitously expressed, 280 kDa type I transmembrane proteoglycan member of the TGF-beta superfamily of proteins. This receptor is a membrane proteoglycan

For Research Use Only

that often functions as a co-receptor with other TGF-beta receptor superfamily members. TGF-beta RIII can enhance or inhibit cell signaling. TGF-beta RIII has been shown to play an essential role in the formation of the atrioventricular cushion and coronary vessels during development of the heart. TGF beta RIII also plays a role in many cancers. Increased expression of TGF beta RIII is found in higher grade lymphomas, and reduced expression of TGF beta RIII is found with advanced stage neuroblastomas and ovarian carcinomas. Low TGF-beta RIII expression also correlates with higher grade among a cohort of breast cancers. TGFBR3 is known to be expressed in adult testis and ovary, but little is known about this receptor during gonadogenesis.