

Recombinant Cynomolgus Thymic Stromal Lymphopoietin/TSLP (C-6His)

Catalog No. PKSQ050102

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Thymic stromal lymphopoietin;Thymic stroma-derived lymphopoietin;TSLP
Species	Cynomolgus macaques
Expression Host	E.coli
Sequence	Tyr29-Gln159(Glu37Gln)
Accession	XP_005557555.1
Calculated Molecular Weight	16.2 kDa
Observed molecular weight	16 kDa
Tag	C-His
Bioactivity	Loaded Anti-Human TSLP mAb-Fc on Protein A Biosensor, can bind Cynomolgus TSLP-His with an affinity constant of 0.07 nM as determined in BLI assay.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 6% Trehalose, 2% Glycine, 50mM NaCl, 0.05% Tween 80, pH7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Thymic stromal lymphopoietin (TSLP) is a protein belonging to the cytokine family, contains 140 amino acids. It is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. TSLP induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c+ dendritic cells. It can induce allergic inflammation by directly activating mast cells. TSLP is produced mainly by non-hematopoietic cells such as fibroblasts, epithelial cells and different types of stromal or stromal-like cells. These cells are located in regions where TSLP activity is required.

For Research Use Only