

Recombinant Human TNFSF4/OX40L protein (His tag)

Catalog No. PDMH100152

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Tumor necrosis factor ligand superfamily member 4;Glycoprotein Gp34;OX40 ligand;OX40L;TAX transcriptionally-activated glycoprotein 1;TNFSF4;CD252;TXGP1;CD134L;CD252;GP34;OX-40;TXGP1
Species	Human
Expression Host	HEK293 Cells
Sequence	Gln51-Leu183
Accession	P23510
Calculated Molecular Weight	14.5 kDa
Observed molecular weight	30 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Tumor necrosis factor ligand superfamily member 4(TNFSF4/OX40L) is a single-pass type II membrane protein. OX40L is expressed on the surface of activated B cells;T cells;dendritic cells and endothelial cells. OX40L binds to OX40(CD134);a member of the TNF receptor superfamily that is expressed predominantly on activated CD4+ T cells. OX40-OX40L co-stimulates signal to promote the survival and proliferation of activated CD4+ T cells and prolong the immune response. It involved in T-cell proliferation and cytokine production. Additional;it has been found association with systemic lupus erythematosus;no association with occurrence of atherosclerosis.

For Research Use Only